

Fixed point conditions for non-coprime actions

Michael C. Burkhart 

University of Cambridge, Cambridge, United Kingdom
(mcb93@cam.ac.uk)

(Received 13 March 2023; accepted 20 August 2023)

In the setting of finite groups, suppose J acts on N via automorphisms so that the induced semidirect product $N \rtimes J$ acts on some non-empty set Ω , with N acting transitively. Glauberman proved that if the orders of J and N are coprime, then J fixes a point in Ω . We consider the non-coprime case and show that if N is abelian and a Sylow p -subgroup of J fixes a point in Ω for each prime p , then J fixes a point in Ω . We also show that if N is nilpotent, $N \rtimes J$ is supersoluble, and a Sylow p -subgroup of J fixes a point in Ω for each prime p , then J fixes a point in Ω .

Keywords: Non-coprime actions; conjugacy of complements; supersoluble groups

2020 *Mathematics Subject Classification* 05E18; 20E22; 20E45; 20F16; 20J06; 55N45

1. Introduction

Suppose a finite group J acts via automorphisms on a finite group N and the induced semi-direct product $G = N \rtimes J$ acts on some non-empty set Ω where the action of N is transitive. Glauberman showed that if each supplement H of N in G splits over $N \cap H$ and each complement of N in G is conjugate to J , then there exists a J -invariant element $\omega \in \Omega$. Consequently, if the orders of J and N are coprime so that the Schur–Zassenhaus theorem applies, a fixed point always exists [4, Thm. 4]. In this note, we consider the non-coprime case and establish some conditions for the existence of a fixed point.

Given an action as described above, consider the stabiliser $G_\alpha \leq G$ fixing an arbitrary point $\alpha \in \Omega$. As N is transitive, G_α supplements N in G . In this context, J fixes an element of Ω if and only if the following two conditions are met. Firstly, we must ensure G_α splits over $N \cap G_\alpha$ so that there exists some complement J' . As $G/N \cong G_\alpha/(N \cap G_\alpha)$, it will follow that J' also complements N in G . Secondly, we require that $J' = g^{-1}Jg$ for some $g \in G$ so that J fixes $g \cdot \alpha$. For the latter requirement, we concern ourselves with conditions for two specific complements in a semidirect product to be conjugate.

To this end, we say two subgroups H and H' are *locally conjugate* in a group G if for each prime p , a Sylow p -subgroup of H is conjugate to a Sylow p -subgroup of H' . Losey and Stonehewer showed that if H and H' are locally conjugate supplements of some normal nilpotent subgroup N in a soluble group G , then H and H' are

conjugate if either G/N is nilpotent or N is abelian [7]. Evans and Shin further showed that if N is abelian, then G need not be soluble [3].

We first restrict N to be abelian and use a decomposition result from group cohomology to provide an alternate proof of:

LEMMA 1.1 (Evans and Shin). *In a finite group, two complements of a normal abelian subgroup are conjugate if and only if they are locally conjugate.*

We use this, along with Gaschütz's result that a finite group G splits over an abelian subgroup N if and only if for each prime p , a Sylow p -subgroup S of G splits over $N \cap S$, to show:

THEOREM 1.2. *Given a finite group J acting via automorphisms on a finite abelian group N , suppose the induced semidirect product $N \rtimes J$ acts on some non-empty set Ω where the action of N is transitive. If for each prime p , a Sylow p -subgroup of J fixes an element of Ω , then there exists some J -invariant element $\omega \in \Omega$.*

This had previously been shown using elementary arguments for the special case that J is supersoluble [2, Cor. 2]. The theorem implies:

COROLLARY 1.3. *Let G be a finite split extension over an abelian subgroup N . If for each prime p there is a Sylow p -subgroup S of G such that any two complements of $N \cap S$ in S are conjugate, then any two complements of N in G are G -conjugate.*

This extends a result of D. G. Higman [5, Cor. 2] that requires the complements of $N \cap S$ in S to be conjugate *within* S .

We then consider nilpotent N and supersoluble $N \rtimes J$. We adapt our approach for lemma 1.1 to nonabelian cohomology and demonstrate:

LEMMA 1.4. *In a finite supersoluble group, two complements of a normal nilpotent subgroup are conjugate if and only if they are locally conjugate.*

With this, we then show:

THEOREM 1.5. *Given a finite group J acting via automorphisms on a finite nilpotent group N , suppose the induced semidirect product $N \rtimes J$ is supersoluble and acts on some non-empty set Ω where the action of N is transitive. If for each prime p , a Sylow p -subgroup of J fixes an element of Ω , then there exists some J -invariant element $\omega \in \Omega$.*

The theorem also implies an analogue of corollary 1.3 that we state and prove in § 3.

1.1. Outline

We proceed as follows. In the remainder of this section, we introduce notation and some conventions from group cohomology. In the next section, we restrict N to be abelian and prove theorem 1.2. We then restrict N to be nilpotent and $N \rtimes J$ to be supersoluble in § 3 and prove theorem 1.5, before concluding in § 4.

1.2. Notation and conventions

All groups in this note are assumed finite. A subgroup $K \leq G$ supplements $N \triangleleft G$ if $G = NK$ and complements N if it both supplements N and the intersection $N \cap K$ is trivial. We denote conjugation by $g^\gamma = \gamma^{-1}g\gamma$ for $g, \gamma \in G$ and otherwise let groups act from the left. For a prime p , we let $\text{Syl}_p(G)$ denote the set of Sylow p -subgroups of a group G .

We rely on rudimentary notions from group cohomology that can be found in the texts of Brown [1] and Serre [8]. Given a group J acting on a group N via automorphisms, crossed homomorphisms or 1-cocycles are maps $\varphi : J \rightarrow N$ satisfying $\varphi(jj') = \varphi(j)\varphi(j')^{j^{-1}}$ for all $j, j' \in J$. Two such maps φ and φ' are cohomologous if there exists $n \in N$ such that $\varphi'(j) = n^{-1}\varphi(j)n^{j^{-1}}$ for all $j \in J$; in this case, we write $\varphi \sim \varphi'$. We take the first cohomology $H^1(J, N)$ to be the pointed set $Z^1(J, N)$ of crossed homomorphisms modulo this equivalence. The distinguished point corresponds to the equivalence class containing the map taking each element of J to the identity of N . Our interest in this set stems primarily from the well-known bijective correspondence [8, Exer. 1 in §I.5.1] between it and the N -conjugacy classes of complements to N in $N \rtimes J$. Specifically, for each $\varphi \in Z^1(J, N)$, the subgroup $F(\varphi) = \{\varphi(j)j\}_{j \in J}$ complements N in NJ and all such complements may be written in this way. Two crossed homomorphisms yield conjugate complements under F if and only if they are cohomologous, so F induces the desired correspondence.

For a subgroup $K \leq J$, we let $\varphi|_K$ denote the restriction of $\varphi \in Z^1(J, N)$ to K and $\text{res}_K^J : H^1(J, N) \rightarrow H^1(K, N)$ be the map induced in cohomology. For $\varphi \in Z^1(K, N)$ and $j \in J$, define $\varphi^j(x) = \varphi(x^{j^{-1}})^j$. We call φ J -invariant if $\text{res}_{K \cap K^j}^K \varphi \sim \text{res}_{K \cap K^j}^{K^j} \varphi^j$ for all $j \in J$ and let $\text{inv}_J H^1(K, N)$ denote the set of J -invariant elements in $H^1(K, N)$. For any $\varphi \in Z^1(J, N)$, we have $\varphi^j(x) = n^{-1}\varphi(x)n^{x^{-1}}$ where $n = \varphi(j^{-1})$ so that $\varphi^j \sim \varphi$. In particular, $\text{res}_K^J H^1(J, N) \subseteq \text{inv}_J H^1(K, N)$.

2. N is abelian

In this section, we restrict N to be abelian so that $H^1(J, N)$ takes the form of an abelian group. We first prove lemma 1.1 as stated in § 1.

Proof of lemma 1.1. Suppose we are given locally conjugate complements J and J' of a normal abelian subgroup N in some group G . As any element $g \in G$ may be uniquely written $g = jn$ for $j \in J$ and $n \in N$, for each prime p we have $J'_p = (J_p)^n$ for some $J_p \in \text{Syl}_p(J)$, $J'_p \in \text{Syl}_p(J')$, and $n \in N$. Let $\varphi' \in Z^1(J, N)$ denote the crossed homomorphism corresponding to J' . It suffices to show that $\varphi' \sim 1$, where $1 \in Z^1(J, N)$ denotes the map taking each element of J to the identity of N . Through the p -primary decomposition of $H^1(J, N)$, we have the isomorphism [1, §III.10]:

$$H^1(J, N) \cong \bigoplus_{p \in \mathcal{D}} \text{inv}_J H^1(J_p, N) \quad (2.1)$$

where \mathcal{D} is the set of prime divisors of $|J|$ and the J_p are those given above. For every $p \in \mathcal{D}$, we see that $\varphi'|_{J_p} \sim 1|_{J_p}$ as J_p and J'_p are N -conjugate complements of N in NJ_p . Thus, φ' maps to the identity in each direct summand on the right-hand side of (2.1) and we may conclude $\varphi' \sim 1$ so that J and J' are conjugate. \square

We can now use the lemma and Gaschütz's theorem to prove theorem 1.2.

Proof of theorem 1.2. Given J , N , and Ω as described in the hypotheses of the theorem, let $G = N \rtimes J$ denote the induced semidirect product and consider the stabiliser subgroup G_α for some fixed $\alpha \in \Omega$. As N acts transitively, any $g \in G$ may be written $g \cdot \alpha = n \cdot \alpha$ for some $n \in N$, so that $n^{-1}g \in G_\alpha$. Thus, $G = NG_\alpha$.

We claim G_α splits over $N \cap G_\alpha$. For any prime p , there exists by hypothesis some $n \in N$ and $P \in \text{Syl}_p(J)$ such that $P^n \leq G_\alpha$. Let $L \in \text{Syl}_p(N \cap G_\alpha)$. As $|G_\alpha| = |N \cap G_\alpha| [G : N]$, it follows that $S = LP^n \in \text{Syl}_p(G_\alpha)$ so P^n complements $S \cap N = L$ in S . As the choice of prime p was arbitrary, we may apply Gaschütz's theorem to conclude that G_α splits over $N \cap G_\alpha$.

Let J' complement $N \cap G_\alpha$ in G_α . As $G/N \cong G_\alpha/(N \cap G_\alpha)$, it follows that J' also complements N in G . Lemma 1.1 then implies that $J' = J^g$ for some $g \in G$ so that J fixes $\omega = g \cdot \alpha$. \square

Finally, we outline how corollary 1.3 follows from theorem 1.2.

Proof of corollary 1.2. Given a group G satisfying the hypotheses of the corollary, suppose J and J' each complement N in G . Then G acts on the cosets $\Omega = G/J'$ in such a way that we may apply theorem 1.2 to infer that J fixes gJ' for some $g \in G$. Therefore, J and J' are conjugate. As the choice of complements was arbitrary, we may conclude. \square

3. N is nilpotent and $N \rtimes J$ is supersoluble

In this section, we suppose that N is nilpotent and $N \rtimes J$ is supersoluble. Consequently, N decomposes as the direct sum $N \cong \oplus_{p \in \mathcal{D}} N_p$ over its characteristic Sylow p -subgroups N_p where \mathcal{D} denotes the set of prime divisors of $|N|$. Direct calculations show that the natural projections $N \rightarrow N_p$ induce an isomorphism of pointed sets

$$H^1(J, N) \cong \oplus_{p \in \mathcal{D}} H^1(J, N_p). \quad (3.1)$$

To parse the components on the right-hand side of (3.1), we introduce the following:

PROPOSITION 3.1. *Suppose a group J acts on a p -group N via automorphisms, so that the induced semidirect product $N \rtimes J$ is supersoluble. Then $\text{res}_{J_p}^J : H^1(J, N) \rightarrow \text{inv}_J H^1(J_p, N)$ is an isomorphism for $J_p \in \text{Syl}_p(J)$.*

Proof. We induct on the order of J . If J itself is a p -group, the conclusion is immediate. If p is not a divisor of $|J|$, the lemma follows from the Schur–Zassenhaus theorem. Otherwise, let $Q \triangleleft J$ be a Sylow q -subgroup where q is the largest prime divisor of $|J|$ [6, Exer. 3B.10] so that $J \cong Q \rtimes M$ for some Hall q' -subgroup $M \leq J$. Consider the inflation–restriction exact sequence [8, §I.5.8],

$$1 \rightarrow H^1(J/Q, N^Q) \rightarrow H^1(J, N) \xrightarrow{\text{res}_Q^J} H^1(Q, N)^{J/Q} \quad (3.2)$$

where N^Q denotes the elements of N fixed by Q .

If $q \neq p$, then $H^1(Q, N)$ is trivial so that $H^1(J, N) \cong H^1(M, N^Q)$. In the supersoluble group NQ , Q is a Sylow q -subgroup for the largest prime divisor of $|NQ|$, so that $Q \triangleleft NQ$ and $N^Q = N$. Consequently, $H^1(J, N) \cong H^1(M, N)$. We claim that res_M^J affords this isomorphism. It suffices to show that res_M^J is surjective. For any $\varphi \in Z^1(M, N)$, we may define $\tilde{\varphi} : J \rightarrow N$ by $\tilde{\varphi}(qm) = \varphi(m)$ for $q \in Q$ and $m \in M$. This map is well-defined as $J \cong Q \rtimes M$. For $q, q' \in Q$ and $m, m' \in M$, we have $\tilde{\varphi}(qmq'm') = \varphi(mm') = \varphi(m)\varphi(m')^{m^{-1}} = \tilde{\varphi}(qm)\tilde{\varphi}(q'm')^{(qm)^{-1}}$, where the last equality follows from the fact that elements of N commute with elements of Q . Thus, $\tilde{\varphi} \in Z^1(J, N)$. As $\tilde{\varphi}|_M = \varphi$, we conclude res_M^J is surjective.

Exchanging M for a conjugate if necessary, we may assume that $J_p \leq M$. As $\text{res}_{J_p}^M$ is injective by induction, it follows that the composition $\text{res}_{J_p}^J = \text{res}_{J_p}^M \circ \text{res}_M^J$ is also injective. On the other hand,

$$\text{inv}_J H^1(J_p, N) \subseteq \text{inv}_M H^1(J_p, N) = \text{res}_{J_p}^M H^1(M, N) \subseteq \text{res}_{J_p}^J H^1(J, N)$$

where the equality above follows from the inductive hypothesis, so that $\text{res}_{J_p}^J$ is surjective.

Otherwise, $q = p$, so that $J_p = Q$ is a Sylow p -subgroup of J . In this case, $H^1(J/Q, N^Q)$ is trivial in (3.2) and so $\text{res}_{J_p}^J$ is injective. As $H^1(Q, N)^{J/Q} = \text{inv}_J H^1(Q, N)$, it remains to show that this map is surjective. For M -invariant $\varphi \in Z^1(J_p, N)$, define $\tilde{\varphi} : J \rightarrow N$ by $\tilde{\varphi}(hm) = \varphi(h)$ for $h \in J_p$ and $m \in M$. Then for any $h, h' \in J_p$ and $m, m' \in M$, we have $\tilde{\varphi}(hmm'h'm') = \varphi(h(h')^{m^{-1}}) = \varphi(h)\varphi((h')^{m^{-1}})^{h^{-1}} = \varphi(h)\varphi(h')^{m^{-1}h^{-1}} = \tilde{\varphi}(hm)\tilde{\varphi}(h'm')^{(hm)^{-1}}$ where the third equality follows from φ being M -invariant. As $J \cong J_p \rtimes M$, we conclude that $\tilde{\varphi} \in Z^1(J, N)$. Clearly, $\text{res}_{J_p}^J \tilde{\varphi} \sim \varphi$ so that $\text{res}_{J_p}^J$ is surjective. \square

For each prime p , we may apply proposition 3.1 to the component for p in (3.1) and find that $H^1(J, N_p) \cong \text{inv}_J H^1(J_p, N_p) \cong \text{inv}_J H^1(J_p, N)$ for some $J_p \in \text{Syl}_p(J)$. In particular, it follows that:

PROPOSITION 3.2. *Given a group J acting on a nilpotent group N via automorphisms so that $N \rtimes J$ is supersoluble, the restriction maps $\text{res}_{J_p}^J$ induce an isomorphism of pointed sets $H^1(J, N) \cong \bigoplus_{p \in \mathcal{D}} \text{inv}_J H^1(J_p, N)$ where \mathcal{D} denotes the set of prime divisors of $|J|$ and $J_p \in \text{Syl}_p(J)$ for each $p \in \mathcal{D}$.*

We are now prepared to provide a proof of lemma 1.4.

Proof of lemma 1.4. In a supersoluble group G , suppose J and J' are locally conjugate complements of a normal nilpotent subgroup N . As in lemma 1.1, we have for each prime p that some $J_p \in \text{Syl}_p(J)$ and $J'_p \in \text{Syl}_p(J')$ are conjugate by an element of N . Let $\varphi' \in Z^1(J, N)$ denote the map corresponding to J' . As the isomorphism in proposition 3.2 is induced by restriction maps, it takes the identity $1 \in H^1(J, N)$ to $\bigoplus_{p \in \mathcal{D}} 1|_{J_p}$. Thus, as $\varphi'|_{J_p} \sim 1|_{J_p}$ for each $p \in \mathcal{D}$, we may apply proposition 3.2 to conclude $\varphi' \sim 1$ so that J and J' are conjugate. \square

We now use lemma 1.4 to show:

PROPOSITION 3.3. *Let H be a subgroup of some supersoluble $G \cong N \rtimes J$ where N is nilpotent. If for each prime p , H contains a conjugate of some $S \in \text{Syl}_p(J)$, then H contains a conjugate of J and so splits over $N \cap H$.*

Proof. The hypotheses imply that H supplements N in G . We induct on the order of G . If N is trivial or if H is a p -group, the conclusion follows immediately. If multiple primes divide $|N|$, then for some prime p , HN_p must be a strict subgroup of G for $N_p \in \text{Syl}_p(N)$; otherwise H would contain a Sylow subgroup of G for each prime and we would have $H = G$. Let p be such a prime. Induction in G/N_p implies $J^g \leq HN_p$ for some $g \in G$. Switching to a conjugate of H if necessary, we may assume that g is trivial and apply the inductive hypothesis in HN_p to conclude $J^{g'} \leq H$ for some $g' \in G$. We now proceed under the assumption that N is a q -subgroup for some prime q .

Let $A \leq N$ be a minimal normal subgroup of G ; as G is supersoluble, it will have prime order q . If $A \leq H$, then in G/A , induction implies that $J^g A \leq HA = H$ for some $g \in G$ so that $J^g \leq H$.

Otherwise, $A \cap H$ is trivial. Without loss, $J_q \leq H$ for some $J_q \in \text{Syl}_q(J)$. In G/A , induction implies that a conjugate of JA/A is contained in HA/A . Let \overline{K} denote this conjugate. Switching to a different conjugate if necessary, we may assume that $J_q A/A \leq \overline{K}$. Let $\varphi: h \mapsto hA/A$ denote the isomorphism from H to HA/A and consider $K = \varphi^{-1}(\overline{K})$. It follows that $J_q \leq K$ and $|K| = |J|$ so that $K \leq H$ complements N in G . As N is a q -group, a Sylow p -subgroup of J will be conjugate to a Sylow p -subgroup of K for primes $p \neq q$. Lemma 1.4 then implies that J and $K \leq H$ are conjugate in G . \square

We now prove theorem 1.5.

Proof of theorem 1.5. Given J , N , and Ω as described in the hypotheses of the theorem, let $G = N \rtimes J$ denote the induced semidirect product and consider G_α for some $\alpha \in \Omega$. As N acts transitively, $G = NG_\alpha$. For each prime p , the hypotheses of the theorem imply $(J_p)^{n_p} \leq G_\alpha$ for some $J_p \in \text{Syl}_p(J)$ and $n_p \in N$, so that proposition 3.3 implies G_α contains a conjugate of J , say J^g for $g \in G$. It follows that J fixes $\omega = g \cdot a$. \square

This in turn implies:

COROLLARY 3.4. *Let G be a supersoluble split extension over a nilpotent subgroup N . If for each prime p there is a Sylow p -subgroup S of G such that any two complements of $S \cap N$ in S are conjugate, then any two complements of N in G are conjugate.*

Proof. Suppose arbitrary J and J' complement N in G . Then G acts on the cosets $\Omega = G/J'$ in such a way that we may apply theorem 1.5 to infer that J fixes gJ' for some $g \in G$. Consequently, J and J' are conjugate, and we may conclude. \square

4. Concluding remarks

In their paper, Losey and Stonehewer exhibited a soluble group $G \cong N \rtimes J$ with N nilpotent and J supersoluble and a second complement J' to N in G such that

J and J' are locally conjugate but not conjugate [7]. Thus, lemma 1.4 cannot be extended to supersoluble complements of a normal nilpotent subgroup in a soluble group.

Acknowledgements

The author thanks Elizabeth Crites, the editor Alex Bartel, and an anonymous reviewer for thoughtful and detailed feedback on the manuscript.

References

- 1 K. S. Brown. *Cohomology of Groups* (Springer, New York, 1982).
- 2 M. C. Burkhart. Conjugacy conditions for supersoluble complements of an abelian base and a fixed point result for non-coprime actions. *Proc. Edinb. Math. Soc. (2)* **65** (2022), 1075–1079.
- 3 M. J. Evans and H. Shin. Local conjugacy in finite groups. *Arch. Math.* **50** (1988), 289–291.
- 4 G. Glauberman. Fixed points in groups with operator groups. *Math. Zeitschr.* **84** (1964), 120–125.
- 5 D. G. Higman. Remarks on splitting extensions. *Pacific J. Math.* **4** (1954), 545–555.
- 6 I. M. Isaacs. *Finite Group Theory* (Amer. Math. Soc., Providence, 2008).
- 7 G. O. Losey and S. E. Stonehewer. Local conjugacy in finite soluble groups. *Quart. J. Math. Oxford (2)* **30** (1979), 183–190.
- 8 J.-P. Serre. *Galois Cohomology* (Springer, Berlin, 2002).